

(D+1)-Dimensional Formulation for D-Dimensional Constrained Systems

Riuji MOCHIZUKI*Department of Physics, Faculty of Science,
Chiba University,
1-33 Yayoi-cho, Chiba 260, Japan*

ABSTRACT

D-dimensional constrained systems are studied with stochastic Lagrangian and Hamiltonian. It is shown that stochastic consistency conditions are second class constraints and Lagrange multiplier fields can be determined in (D+1)-dimensional canonical formulation. The Langevin equations for the constrained system are obtained as Hamilton's equations of motion where conjugate momenta play a part of noise fields.

The stochastic quantization scheme was first proposed by Parisi and Wu^[1] as an alternative quantization method for gauge theories^[2] in 1981. After a few years Namiki *et al.*^[3] improved the method so as to handle constrained systems and recently more rigorous discussion was shown.^[4] In their methods Lagrange multiplier fields can be determined under stochastic consistency conditions (the consistency conditions for the fictitious time). If we extend the stochastic quantization method to phase space, we can determine the Lagrange multipliers accompanied with the first class constraints by almost the same method.^[5] They seem strange because we cannot determine them in canonical formalism. In this letter we introduce the stochastic Lagrangian^[6] and Hamiltonian for constrained systems and treat them within a (D+1)-dimensional canonical framework. It is shown that stochastic consistency conditions are second class constraints and we obtain the Langevin equations for the constrained system as Hamilton's equations of motion.

We consider a system described by the action S in D-dimensional Euclidean space-time. In the stochastic quantization scheme the Langevin equation is an important tool, which is written in the discretized form as

$$dq_i = -\frac{\delta S}{\delta q_i}dt + dW_i. \quad (i = 1, \dots, N) \quad (1)$$

Here t is the fictitious time and dW_i is the Wiener process whose expectation values are defined as

$$\langle dW_i(t)dW_j(t') \dots \rangle = \frac{1}{n} \int D(dW) dW_i(t)dW_j(t') \dots \exp \left\{ - \int dx \sum_m L(dW(\tau_m)) \right\}, \quad (2)$$

where n is a normalization factor and $L(dW)$ is stochastic Lagrangian. If the system has constraints

$$F_a = 0, \quad (a = 1, \dots, M; \ N > M) \quad (3)$$

we define the stochastic Lagrangian as

$$L(dW) \equiv \frac{1}{4dt} dW_i dW_i + \lambda_a dF_a, \quad (4)$$

where λ_a is a Lagrange multiplier field. In the stochastic Lagrangian (4) we have changed the constraints (3) into their differentials. It is quite natural in this formulation because the order of the stochastic Lagrangian is dt and we rewrite it with the Langevin equations, i.e. stochastic differential equations, as shown in the following.

When we handle stochastic differential equations, we should make choice of their calculation rules, that is, adopt either Ito's calculation rule^[7] or Stratonovich's one^[8] in a general way. If we adopt the latter rule, we must bother about the Jacobian factor which appears when we change the integration variables in the generating functional. On the other hand the Jacobian factor is a unity if we calculate with Ito's rule,^[6] therefore we will use it in the following calculation. In the Ito calculus we have Ito's formula

$$\langle df \rangle = \left\langle \frac{\partial f}{\partial q_i} dq_i \right\rangle + \frac{1}{2} \left\langle \frac{\partial^2 f}{\partial q_i \partial q_j} dW_i dW_j \right\rangle \quad (5)$$

where f is an arbitrary functional of q_i 's.

To change the integration variables we insert a unity

$$1 = \int D(q) \delta\{dq_i + \frac{\delta S}{\delta q_i} dt - dW_i\}$$

into RHS of (2) and integrate over dW . Then the distribution functional Z reads

$$Z = \int D(q) \exp\left\{-\int dx \sum L(q)\right\}, \quad (6)$$

where

$$L(q) = \frac{1}{4dt} (dq_i + \frac{\delta S}{\delta q_i} dt)^2 + \lambda_a dF_a, \quad (7)$$

$$dF_a = \frac{\partial F_a}{\partial q_i} dq_i + \frac{1}{2} \cdot \frac{\partial^2 F_a}{\partial q_i \partial q_j} M_{ij}. \quad (8)$$

The second term in RHS of (8) is due to Ito's formula, M_{ij} in which is a dt -order expectation value of $dW_i dW_j$ which is not $2\delta_{ij}$ because the Lagrange multiplier fields should contain a noise-like part.^[4] We will compute it later.

From the stochastic Lagrangian (7) we define stochastic conjugate momenta as

$$p_i \equiv \frac{\partial L}{\partial(dq_i)} = \frac{1}{2} \left(\frac{dq_i}{dt} + \frac{\partial L}{\partial q_i} \right) + \lambda_a \frac{\partial F_a}{\partial q_i}, \quad (9.a)$$

$$\pi_a \equiv \frac{\partial L}{\partial(d\lambda_a)} \approx 0. \quad (9.b)$$

The latter equation is a primary constraint, and hence the stochastic Hamiltonian H is

$$\begin{aligned} H &\equiv p_i dq_i + \pi_a d\lambda_a - L + u_a \pi_a dt \\ &= p_i (2p_i dt - \frac{\delta S}{\delta q_i} dt - 2\lambda_a \frac{\partial F_a}{\partial q_i} dt) - \frac{1}{dt} (p_i dt - \lambda_a \frac{\partial F_a}{\partial q_i} dt) (p_i dt - \lambda_b \frac{\partial F_b}{\partial q_i} dt) \\ &\quad - \lambda_a \left\{ \frac{\partial F_a}{\partial q_i} (2p_i dt - \frac{\delta S}{\delta q_i} dt - 2\lambda_b \frac{\partial F_b}{\partial q_i} dt) + \frac{1}{2} \cdot \frac{\partial^2 F_a}{\partial q_i \partial q_j} M_{ij} \right\} + u_a \pi_a dt, \end{aligned} \quad (10)$$

where u_a is the other Lagrange multiplier into which $d\lambda_a/dt$ is absorbed.

The stochastic differentials are defined using Poisson's bracket

$$\{A, B\} \equiv \frac{\partial A}{\partial q_i} \frac{\partial B}{\partial p_i} - \frac{\partial A}{\partial p_i} \frac{\partial B}{\partial q_i} + \frac{\partial A}{\partial \lambda_a} \frac{\partial B}{\partial \pi_a} - \frac{\partial A}{\partial \pi_a} \frac{\partial B}{\partial \lambda_a}$$

as

$$dA \equiv \{A, H\}. \quad (11)$$

The stochastic consistency condition^[3] of $\pi_a \approx 0$ thus reads

$$\begin{aligned} d\pi_a &= \{\pi_a, H\} \\ &= \frac{\partial F_a}{\partial q_i} (2p_i dt - \frac{\delta S}{\delta q_i} dt) + \frac{1}{2} \cdot \frac{\partial^2 F_a}{\partial q_i \partial q_j} M_{ij} - 2 \frac{\partial F_a}{\partial q_i} \frac{\partial F_b}{\partial q_i} \lambda_b dt \\ &\equiv \phi_a \approx 0. \end{aligned} \quad (12)$$

This is a secondary constraint. If we determine λ_b so as to satisfy it, we need no more constraints. We may identify ϕ_a with dF_a and π_a with $F_a + C_a$ where C_a is an arbitrary functional independent of the fictitious time. Consequently we should choose the initial conditions as $C_a = 0$ to realize the constraints (3). The Poisson's bracket of the two constraints (9.b) and (12) is

$$\{\pi_a, \phi_b\} = 2 \frac{\partial F_a}{\partial q_i} \frac{\partial F_b}{\partial q_i} \equiv 2D_{ab}. \quad (13)$$

Here we assume that D_{ab} is not zero and has its inverse D_{ab}^{-1} ^[3]. Then the above equation shows that both of them are second class constraints. When we write the total Hamiltonian H_T as

$$H_T = H + v_a \pi_a dt + w_a \phi_a \quad (14)$$

where v_a and w_a are Lagrange multiplier fields, we can determine them with the help of the stochastic consistency conditions again. They read

$$w_a = 0,$$

$$v_a = \frac{1}{2} D_{ab}^{-1} \{\phi_b, H\}.$$

Accordingly the total Hamiltonian becomes

$$H_T = H + \frac{1}{2} D_{ab}^{-1} \{\phi_b, H\} \pi_a. \quad (15)$$

Next we derive Hamilton's equation of motion for q_i . If we use Dirac's bracket

$$[A, B] \equiv \{A, B\} - \{A, \Phi_r\} \Delta_{rs}^{-1} \{\Phi_s, B\} \quad (16)$$

with

$$\Phi_r \equiv \pi_a, \phi_a,$$

$$\Delta_{rs} \equiv \{\Phi_r, \Phi_s\},$$

we may replace the weak equality $\approx$ by the strong equality $=$. In the following we always use Dirac's bracket instead of Poisson's bracket and rewrite the total Hamiltonian by the aid of the second class constraints (9.b) and (12) as

$$\begin{aligned} H_T = & p_i p_i dt - p_i \frac{\delta S}{\delta q_i} dt \\ & - \frac{1}{4dt} D_{ab}^{-1} \left\{ \frac{\partial F_a}{\partial q_i} (2p_i dt - \frac{\delta S}{\delta q_i} dt) + \frac{1}{2} \cdot \frac{\partial^2 F_a}{\partial q_i \partial q_k} M_{ik} \right\} \\ & \times \left\{ \frac{\partial F_b}{\partial q_j} (2p_j dt - \frac{\delta S}{\delta q_j} dt) + \frac{1}{2} \cdot \frac{\partial^2 F_b}{\partial q_j \partial q_l} M_{jl} \right\}. \end{aligned} \quad (17)$$

Using it we obtain Hamilton's equation of motion for q_i :

$$\begin{aligned} dq_i = & [q_i, H_T] \\ = & -K_{ij} \frac{\delta S}{\delta q_j} dt + 2K_{ij} p_j dt - \frac{1}{2} \cdot \frac{\partial F_a}{\partial q_i} D_{ab}^{-1} \frac{\partial^2 F_b}{\partial q_j \partial q_k} M_{jk} \end{aligned} \quad (18)$$

with

$$K_{ij} \equiv \delta_{ij} - \frac{\partial F_a}{\partial q_i} D_{ab}^{-1} \frac{\partial F_b}{\partial q_j},$$

which of course satisfies $dF_a = 0$, and hence $(N - M)$ independent equations exist.

The equation (18) is Langevin equation if we may identify $2K_{ij} p_j dt$ with noise fields. In the following we show that is the case. In order to decompose the variables into constraint variables and independent ones,^{[3][4]} we introduce a new set of variables $\{Q^\mu\}$, ($\mu =$

$1, 2, \dots, N$), which are defined as

$$\frac{\partial Q^\mu}{\partial q_i} = e^\mu_i, \quad (19)$$

where e^μ_i 's are vielbein fields defined as

$$e^a_i \equiv \frac{\partial F_a}{\partial q_i}, \quad e_{a,i} \equiv D_{ab}^{-1} \frac{\partial F_b}{\partial q_i}, \quad (a = N - M + 1, \dots, N) \quad (20.a)$$

$$e^A_i e_{a,i} = 0, \quad (A = 1, 2, \dots, N - M) \quad (20.b)$$

$$e_{A,i} \equiv (g^{-1})_{AB} e^B_i, \quad g^{AB} \equiv e^A_i e^B_i. \quad (20.c)$$

Here we have assumed that the metric g^{AB} is non-singular. These definitions lead to following relations

$$e^\mu_i e_{\nu,i} = \delta_{\mu\nu}, \quad e^\mu_i e_{\mu,j} = \delta_{ij}, \quad (21.a)$$

$$e^A_i e^a_i = e^A_i e_{a,i} = e^{A,i} e^a_i k = e_{A,i} e_{a,i} = 0, \quad (21.b)$$

$$e^A_i e_{A,j} = K_{ij}, \quad e^a_i e_{a,j} = \delta_{ij} - K_{ij}, \quad (21.c)$$

$$K_{ij} e^a_j = K_{ij} e_{a,j} = 0, \quad (\delta_{ij} - K_{ij}) e^A_j = (\delta_{ij} - K_{ij}) e_{A,j} = 0, \quad (21.d)$$

$$\det(e^\mu_i) \neq 0. \quad (21.e)$$

The equations (21.e) shows that a manifold spanned by Q^μ 's is equivalent to one by q_i 's.

Assuming $2K_{ij}p_j dt$ to be the noise field and to give the same expectation value as dW_i , which assumption will be confirmed later, we can derive the Langevin equations for the new variables. Taking account of Ito's formula (5), they are

$$dQ^\mu = e^\mu_i dq_i + \frac{1}{2} \frac{\partial e^\mu_i}{\partial q_j} M_{ij}. \quad (22)$$

For the constraint variable $Q^a = F_a$, we have

$$dQ^a = 0 \quad (23.a)$$

and the initial conditions should be chosen to be zero to realize the constraints (3),. On the

other hand, for the independent variables $Q^{A'}$ s, the Langevin equations are

$$dQ^A = -g^{AB} \frac{\delta S}{\delta Q^B} dt + e_i^A (2p_i dt) + \frac{1}{2} \frac{\partial e_i^A}{\partial Q^B} e_j^B M_{ij}. \quad (23.b)$$

Moreover if we decompose the vielbein e_i^A as

$$e_i^A \equiv E_I^A \epsilon_i^I, \quad (I = 1, 2, \dots, N - M) \quad (24.a)$$

$$\epsilon_i^I \epsilon_j^J = \delta^{IJ}, \quad e_i^A e_j^B = E_I^A E_J^B \delta^{IJ} = g^{AB}, \quad (24.b)$$

where ϵ_i^I only rotates the tangent space spanned by the independent variables, we may change the second term in RHS of the Langevin equation (23.b) into fields with $N - M$ indices:

$$e_i^A (2p_i dt) = E_I^A d\Omega^I, \quad (25)$$

where $d\Omega^I$ is defined as

$$d\Omega^I \equiv \epsilon_i^I (2p_i dt), \quad (26)$$

which will be regarded as the Wiener process as shown in the following.

We change the variables q_i 's in the distribution functional (6) into the new ones Q^μ 's

$$Z = \int D(Q) \det(e_{\mu,i}) \exp \left\{ - \int dx \sum L(Q) \right\}, \quad (27)$$

and substitute zero for Q^a 's. Then again we insert a unity

$$1 = \int D(d\Omega) \det(E_I^A) \delta \left\{ dQ^A + g^{AB} \frac{\delta S}{\delta Q^B} dt - E_I^A d\Omega^I - \frac{1}{2} \frac{\partial e_i^A}{\partial Q^B} e_j^B M_{ij} \right\}$$

into the distribution functional (27) and integrate over Q^A 's. Neglecting the normalization

factor, (27) becomes

$$Z = \int D(d\Omega) \exp\left\{-\int dx \sum L(d\Omega)\right\}, \quad (28)$$

where

$$L(d\Omega) = \frac{1}{4dt} d\Omega_I d\Omega_I. \quad (29)$$

The expectation values are easily computed with the above distribution functional:

$$\langle d\Omega^I \rangle = 0,$$

$$\langle d\Omega^I d\Omega^J \rangle = 2\delta^{IJ} dt,$$

and

$$M_{ij} = 2\langle e^A_i e_{A,j} \rangle dt.$$

The Langevin equation (23.b) thus becomes

$$dQ^A = -g^{AB} \frac{\delta S}{\delta Q^B} dt + E^A_I d\Omega^I + \frac{\partial e^A_i}{\partial Q^B} e^B_i dt, \quad (30)$$

which agrees with the Langevin equation for constrained systems obtained by the other method in the stochastic quantization scheme.^[4] The last term is necessary for the general coordinate transformation covariance^[9] and, in perturbation theories, cancellation of some of divergent terms.^[4] The proper Fokker-Planck equation on $(N - M)$ -dimensional constraint surface is given by the ordinary prescription with the Langevin equations (23). It will be straightforward to extend the results to the phase-space stochastic quantization scheme.^[5]

Finally we comment on the results when we obey Stratonovich's calculation rule. In this case the Langevin equation (30) should be a little modified.^{[4][10][11]} Nevertheless it, of course, does not mean the two rules lead different physical results. For example the Fokker-Planck equation, whose equilibrium distribution should be consistent with the path-integral form, is the same in both the cases.

1. ACKNOWLEDGMENTS

I would like to thank Professor T.Kimura, Dr. A.Nakamura and K.Ikegami for useful discussions.

REFERENCES

1. G.Parisi and Y.Wu, Sci.Sin. **24** (1981) 483
2. For a review, P.H.Damgaard and H.Hüffel, Phys.Rep. **152** (1987) 227
3. M.Namiki, I.Ohba and K.Okano, Prog.Theor.Phys. **72** (1984) 350
4. K.Ikegami, T.Kimura and R.Mochizuki, CHIBA-EP-57 (1992)
5. R.Mochizuki, CHIBA-EP-56 (1991)
6. M.Namiki and Y.Yamanaka, Prog.Theor.Phys. **69** (1983) 1764,
E.Gozzi, Phys.Rev. **D28** (1983) 1922,
Y.Nakano, Prog.Theor.Phys. **69** (1983) 361,
C.M.Bender, F.Cooper and B.Freedman, Nucl.Phys. **B219** (1983) 61
7. K.Ito, Proc.Imp.Acad. **20** (1944) 519
8. R.L.Stratonovich, *Conditional Markov Processes and Their Application to the Theory of Optimal Control* (Elsevier, New York, 1968)
9. R.Graham, Phys.Lett. **109A** (1985) 209
10. J.Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Clarendon Press, Oxford, 1989),
R.Mochizuki, Mod.Phys.Lett. **A5** (1990) 2335
11. R.Mochizuki, Prog.Theor.Phys. **85** (1991) 407